

$$\gamma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \gamma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \gamma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

where I_n is the identity matrix of dimension n . Solutions to Eq. (1) are complex-valued 4-component functions and can describe both electronic (positive energy) and positronic (negative energy) states. Methods exist to transform the Dirac equation such that one must only solve a two-component problem for the electronic states. Good reviews of these methods exist, for example, Refs. 10 and 11. Although it is possible to use MADNESS in the context of a two-component formalism, the computational cost is prohibitive for practical calculations as discussed in Ref. 12. Therefore, in this paper, we concern ourselves only with the full 4-component Hamiltonian.

The adaptive multiwavelet basis (to be described in Sec. I B) in MADNESS is not band-limited, so solving Eq. (1) by forming the equivalent matrix eigenvalue problem in the space of basis functions would amplify high frequency noise in the calculation of the matrix elements. For this reason, the integral equation is solved instead by using the associated Green's function. This method has been successfully applied in the nonrelativistic case,¹³ and we extend it to the relativistic case here. Rearranging (1) and application of the matrix Green's function associated with the free-particle Dirac equation gives

$$(\hbar_d - i) \psi(\mathbf{r}) = (K - J - V) \psi(\mathbf{r}), \quad (5)$$

$$\psi = G_i(\mathbf{r}) [(K - J - V) \psi(\mathbf{r})]. \quad (6)$$

Above, the ψ represents convolution of the two operands. G_i is the matrix Green's function for the operator on the left-hand side of Eq. (5), i.e., G_i solves

$$(\hbar_d - i) G_i(\mathbf{r}) = \delta(\mathbf{r}) I_4. \quad (7)$$

Its form, provided with proof in Ref. 14, is

$$G_i(\mathbf{r}) = \frac{1}{2mc^2} [\hbar_d + i] g_i(\mathbf{r}), \quad (8)$$

$$g_i(\mathbf{r}) = \frac{e^{-i\mathbf{r}}}{4|\mathbf{r}|}, \quad (9)$$

$$i = \frac{m^2 c^4 - \frac{1}{2}}{mc^2}. \quad (10)$$

Note that the form of Eqs. (8)–(10) changes if the relativistic Hamiltonian used subtracts the rest energy from the standard

II. METHODS

Equation (6) leads to a method of iterating a function represented in the multiwavelet basis to convergence. Letting n represent the calculation iteration, we have

$$r_i^{n+1} = G_i^n(r) \quad [(K - J -$$

TABLE I. Comparison of MADNESS and GRASP Dirac-Fock ground state energy calculations for atoms.

Atom	MADNESS	GRASP	Error	Relative error
Be	-14.575 89	-14.575 89	-9.7×10^{-7}	6.7×10^{-8}
Ne	-128.691 94	-128.691 93	-8.9×10^{-6}	6.9×10^{-8}
Mg	-199.935 09	-199.935 07	-2.1×10^{-5}	1.1×10^{-7}
Ar	-528.683 80	-528.683 76	-3.6×10^{-5}	6.8×10^{-8}
Ca	-679.710 20	-679.710 16	-3.9×10^{-5}	5.7×10^{-8}
Zn	-1 794.613 01	-1 794.612 97	-3.1×10^{-5}	1.7×10^{-8}
Kr	-2 788.860 65	-2 788.860 58	6.8×10^{-5}	2.4×10^{-8}
Sr	-3 178.079 90	-3 178.079 91	8.1×10^{-6}	-2.5×10^{-9}
Cd	--			

to GRASP and DIRAC. While this approach is unlikely to replace grid-based calculations for atoms in the near future, to the best of our knowledge it is the only fully numerical approach for general molecules.

ACKNOWLEDGMENTS

J. Anderson, R. Harrison, and B. Sundahl were supported in part by the NSF under Grant No. ACI-1450344. The authors would like to acknowledge Dr. William Scott Thornton, Dr. Hideo Sekino, Dr. Bhanu Das, and Dr. Srinivasa Prasanna for helpful discussions on relativistic quantum chemistry theory and current numerical methods.

REFERENCES

- ¹P. Pyykkö, "Relativistic effects in structural chemistry," *Chem. Rev.* **88**, 563–594 (1988).
- ²D. Moncrieff and S. Wilson, "Computational linear dependence in molecular electronic structure calculations using universal basis sets," *Int. J. Quantum Chem.* **101**, 363–371 (2005).
- ³R. E. Stanton and S. Havriliak, "Kinetic balance: A partial solution to the problem of variational safety in Dirac calculations," *J. Chem. Phys.* **81**, 1910 (1984).
- ⁴F. Parpia and I. Grant, "Software for relativistic atomic theory: The GRASP project at Oxford," *J. Phys. IV Colloque* **01**, 33–46 (1991).
- ⁵A. Rosén and D. Ellis, "Relativistic molecular wavefunctions: XeF₂," *Chem. Phys. Lett.* **27**, 595–599 (1974).
- ⁶J. Desclaux and P. Pyykkö, "Relativistic and non-relativistic Hartree-Fock one-centre expansion calculations for the series CH₄ to PbH₄ within the spherical approximation," *Chem. Phys. Lett.* **29**, 534–539 (1974).
- ⁷D. Sundholm, P. Pyykkö, and L. Laaksonen, "Two-dimensional, fully numerical solutions of second-order dirac equations for diatomic molecules. Part 3," *Phys. Scr.* **36**, 400–402 (1987).
- ⁸R. J. Harrison, G. Beylkin, F. A. Bischoff, J. A. Calvin, G. I. Fann, J. Fosso-tande, D. Galindo, J. R. Hammond, R. Hartman-baker, J. C. Hill, J. Jia, J. S. Jakob, M.-J. Yvonne Ou, J. Pei, L. E. Ratcliff, M. G. Reuter, A. C. Richie-Halford, N. A. Romero, H. Sekino, W. A. Shelton, B. E. Sundahl, W. Scott, E. F. Valeev, A. Vazquez-Mayagoitia, N. Vence, T. Yanai, and Y. Yokoi, "Madness: A multiresolution, adaptive numerical environment for scientific simulation," *SIAM J. Sci. Comput.* **38**, S123–S142 (2016).
- ⁹L. Visscher and K. Dyall, "Dirac-Fock atomic electronic structure calculations using different nuclear charge distributions," *At. Data Nucl. Data Tables* **67**, 207–224 (1997).
- ¹⁰W. Liu, "Ideas of relativistic quantum chemistry," *Mol. Phys.* **108**, 1679–1706 (2010).
- ¹¹T. Saue, "Relativistic Hamiltonians for chemistry," *AIP Conf. Proc.* **1504**, 219–227 (2012).
- ¹²J. Anderson, R. Harrison, B. Sundahl, W. Thornton, and G. Beylkin, "Real-space quasi-relativistic quantum chemistry," *Computational and Theoretical Chemistry* (submitted).
- ¹³R. J. Harrison, G. I. Fann, T. Yanai, Z. Gan, and G. Beylkin, "Multiresolution quantum chemistry: Basic theory and initial applications," *J. Chem. Phys.* **121**, 11587–11598 (2004).
- ¹⁴J. Blackledge and B. Babajanov, "On the Dirac scattering problem," *Math. Aeterna* **3**, 535–544 (2013).
- ¹⁵B. Alpert, G. Beylkin, D. Gines, and L. Vozovoi, "Adaptive solution of partial differential equations in multiwavelet bases," *J. Comput. Phys.* **182**, 149–190 (2002).
- ¹⁶H. M. Kalos, "Monte Carlo calculations of the ground state of three- and four-body nuclei," *Phys. Rev.* **128**, 1791–1795 (1962).
- ¹⁷M. Valiev, E. J. Bylaska, N. Govind, K. Kowalski, T. P. Straatsma, H. J. J. Van Dam, D. Wang, J. Nieplocha, E. Apra, T. L. Windus, and W. A. De Jong, "NWChem: A comprehensive and scalable open-source solution for large scale molecular simulations," *Comput. Phys. Commun.* **181**, 1477–1489 (2010).
- ¹⁸R. J. Harrison, "Krylov subspace accelerated inexact Newton method for linear and nonlinear equations," *J. Comput. Chem.* **25**, 328–334 (2004).
- ¹⁹DIRAC, a relativistic *ab initio* electronic structure program, release DIRAC15 (2015), written by R. Bast, T. Saue, L. Visscher, and H. J. Aa. Jensen, with contributions from V. Bakken, K. G. Dyall, S. Dubillard, U. Ekstroem, E. Eliav, T. Enevoldsen, E. Fasshauer, T. Fleig, O. Fossgaard, A. S. P. Gomes, T. Helgaker, J. Henriksson, M. Ilias, Ch. R. Jacob, S. Knecht, S. Komorovsky, O. Kullie, J. K. Laerdahl, C. V. Larsen, Y. S. Lee, H. S. Nataraj, M. K. Nayak, P. Norman, G. Olejniczak, J. Olsen, Y. C. Park, J. K. Pedersen, M. Pernpointner, R. Di Remigio, K. Ruud, P. Salek, B. Schimmelpfennig, J. Sikkema, A. J. Thorvaldsen, J. Thyssen, J. van Stralen, S. Villaume, O. Visser, T. Winther, and S. Yamamoto, see <http://www.diracprogram.org>.
- ²⁰K. G. Dyall, *Theor. Chem. Acc.* **99**, 366 (1995 Td625(Chem.)-625(Acc.))TJ0 g 0 G/F/