

1. (20 pts) Parts (a) and (b) are not related.

- (a) For $f(x) = \frac{1}{x^2}$ and $g(x) = \frac{1}{x+2}$, identify the composite function $(f \circ g)(x)$ and its domain. Express the domain in interval form.

Solution:


$$(f \circ g)(x) = f(g(x)) = f\left(\frac{1}{x+2}\right) = \frac{1}{\left(\frac{1}{x+2}\right)^2} = (x+2)^2 = \boxed{x+2}$$

The domain of $f \circ g$ is the set of all x in the domain of g such that $g(x)$ is in the domain of f .

$$\text{Domain of } g: \left(x+2 > 0 \right) \Rightarrow x > -2$$


For each x in the interval $(-2; \infty)$, $g(x)$ is in the domain of f (since $g(x) \neq 0$ for all x).

- (b) The graph below depicts a function of the form $y = h(x) = a \sin (bx) + c$. Determine the values of a , b , and c . (*Hint: Consider the transformations from the graph of $y = \sin x$ to the given graph.*)



Solution:

Begin with the graph of the relevant base curve, $y = \sin x$:



The profile of the given curve over the interval $[0; 12]$ is the same as the profile of the $y = \sin x$ curve over the interval $[0; 3]$. Therefore, the given curve has experienced a horizontal compression of a factor of 3 with respect to the $y = \sin x$ curve, which implies that $b = 3$

The vertical difference between the given curve's maximum and minimum values is $1 - (-1) = 2$, while the vertical difference between the $y = \sin x$ curve's maximum and minimum values is $1 - (-1) = 2$. Therefore, the given curve has experienced a vertical expansion of a factor of 2 with respect to the $y = \sin x$ curve, which implies that $a = 2$

The vertical center of the given curve is $y = -1$ while the vertical center of the $y = \sin x$ curve is $y = 0$. Therefore, the given curve has experienced a downward vertical shift of 1 unit with respect to the $y = \sin x$ curve, which implies that $c = -1$

Therefore, the function depicted in the given graph is $y = 2 \sin (3x) - 1$

2. (30 pts) Evaluate the following limits. Support your answers by stating theorems, definitions, or other key properties that are used.

(a) $\lim_{x \neq 0} \frac{\tan x \sin(2x)}{x^2}$

Solution: Key property: $\lim_{x \neq 0} \frac{\sin x}{x} = 1$

$$\begin{aligned} \lim_{x \neq 0} \frac{\tan x \sin(2x)}{x^2} &= \lim_{x \neq 0} \frac{\tan x}{x} \cdot \frac{\sin(2x)}{x} \\ &= \lim_{x \neq 0} \frac{\sin x}{x \cos x} \cdot \frac{2 \sin(2x)}{2x} \\ &= \lim_{x \neq 0} \frac{\sin x}{x} \cdot \frac{1}{\cos x} \cdot \frac{2 \sin(2x)}{2x} \\ &= \lim_{x \neq 0} \frac{\sin x}{x} \cdot \lim_{x \neq 0} \frac{2}{\cos x} \cdot \lim_{x \neq 0} \frac{\sin(2x)}{2x} \\ &= [1] \cdot \frac{2}{1} \end{aligned}$$

$$(b) \lim_{x \rightarrow 9} \frac{\sqrt{x-5} - 2}{x-9}$$

Solution:

Begin by multiplying the numerator and the denominator by the conjugate of the original numerator expression.

$$\lim_{x \rightarrow 9} \frac{\sqrt{x-5} - 2}{x-9} = \lim_{x \rightarrow 9} \frac{\sqrt{x-5} - 2}{x-9} \cdot \frac{\sqrt{x-5} + 2}{\sqrt{x-5} + 2}$$

3. (30 pts) Consider the rational function $r(x) = \frac{x^2 - 5x + 4}{2x^2 - 8x + 6}$.

- (a) Identify all values of x at which $r(x)$ is discontinuous. At each such x value, explain why the function is discontinuous there.

Solution:

$$r(x) = \frac{x^2 - 5x + 4}{2x^2 - 8x + 6} = \frac{(x - 1)(x - 4)}{2(x - 1)(x - 3)}$$

Since $r(x)$ is a rational function, it is continuous at all x in its domain.

- (c) Find the equation of each vertical asymptote of $y = r(x)$, if any exist. Support your answer in terms of the limits you evaluated in part (b).

Solution:

The finite value of $\lim_{x \rightarrow 1} r(x) = \frac{3}{4}$ determined in part (b) indicates that there is no vertical asymptote at $x = 1$.

The infinite limits $\lim_{x \rightarrow 3} r(x) = 1$ and $\lim_{x \rightarrow 3^+} r(x) = 1$ were determined in part (b). Either one of those limits being infinite is sufficient to conclude that the line $x = 3$ is a vertical asymptote of the curve $y = r(x)$.

- (d) Find the equation of each horizontal asymptote of $y = r(x)$, if any exist. Support your answer by evaluating the appropriate limits.

Solution:

$$\begin{aligned}\lim_{x \rightarrow 1} r(x) &= \lim_{x \rightarrow 1} \frac{x^2 - 5x + 4}{2x^2 - 8x + 6} = \lim_{x \rightarrow 1} \frac{x^2 - 5x + 4}{2x^2 - 8x + 6} \cdot \frac{1-x^2}{1-x^2} \\ &= \lim_{x \rightarrow 1} \frac{1 - 5 + 4 - x^2}{2 - 8 + 6 - x^2} = \frac{1 - 0 + 0}{2 - 0 + 0} = \frac{1}{2}\end{aligned}$$

Therefore, the equation of the only horizontal asymptote is $y = \frac{1}{2}$.

4. (20 pts) Parts (a) and (b) are not related.

- (a) For what value of b is the following function $u(x)$ continuous at $x = 3$? Support your answer using the definition of continuity, which includes evaluating the appropriate limits.

$$u(x) = \begin{cases} \frac{x^2 - 9}{x - 3} & ; x < 3 \\ 5x + b & ; x \geq 3 \end{cases}$$

Solution:

By the definition of continuity, $u(x)$ is continuous at $x = 3$ if $\lim_{x \rightarrow 3^-} u(x) = \lim_{x \rightarrow 3^+} u(x) = u(3)$.

$$\lim_{x \rightarrow 3^-} u(x) = \lim_{x \rightarrow 3^-} \frac{x^2 - 9}{x - 3} = \lim_{x \rightarrow 3^-} \frac{(x - 3)(x + 3)}{x - 3} = \lim_{x \rightarrow 3^-} (x + 3) = 3 + 3 = 6$$

$$\lim_{x \rightarrow 3^+} u(x) = \lim_{x \rightarrow 3^+} (5x + b) = (5)(3) + b = 15 + b$$

$$u(3) = (5)(3) + b = 15 + b$$

Therefore, $u(x)$ is continuous at $x = 3$ if $6 = 15 + b$, which occurs when $b = -9$

- (b) The Intermediate Value Theorem can **NOT** be used to guarantee that $v(x) = \frac{2}{x} + \frac{1}{x+2} = 0$ for a value of x on the interval $(-1; 2)$. Explain which condition for applying the theorem is not satisfied in this case.

Solution:

The Intermediate Value Theorem cannot be applied in this case because $v(0)$ is undefined, which means that

$v(x)$ is not continuous on the interval $[-1; 2]$

The continuity of $v(x)$ on $[-1; 2]$ is one of the hypotheses for applying the IVT to the given function on the given interval.

(Note that $v(-1) = -1$ and $v(2) = 3$ together indicate that the other IVT hypothesis does hold)