



Iterated Spherical Means in Linearized Inverse Problems

Gregory Beylkin*

Abstract. We consider a representation of the function

$$f(x) = \int_{S^{n-1}} \hat{F}(p) dp$$

in terms of the iterated spherical mean of $\hat{F}(p)$. Here, n is the dimension of the space. We also review applications of such a representation to linearized inverse problems and present

$2\pi^{n/2}$

is a point in D^n is a

If the support of $\hat{F}(p)$ is not restricted to the ball B_{2k} in (3.2), then (3.4) defines the low-pass-filtered version of the function whose Fourier transform is $\hat{F}(p)$. In this case (3.3) is the representation of the low-pass-filtered version of the function in terms of the iterated spherical mean of its Fourier transform.

The representation in (3.3) was derived in [4]. For $n = 2$ and $n = 3$ it reduces to formulae obtained by A. J. Devaney [1,3].

4. Inverse Scattering in Born Approximations. The most simple example of an application of the representation in (3.3) is the inversion formula for inverse scattering

and

$$Q_{LP}(x) = \frac{k^3}{\pi^2} \int_{L^+} \int_{L^+} |\nu - \mu| |f(k, \nu, \mu)|^2 e^{i(k\nu - k\mu) \cdot x} d\omega_\nu d\omega_\mu. \quad (4.6)$$

Thus, we obtain that if we can measure the phase of the scattering amplitude we have the explicit inversion formula in (4.5) for the reconstruction of the low-pass-filtered version of the potential. In the case when the phase of the scattering amplitude cannot be directly measured we can explicitly reconstruct the interatomic distance function using (4.6). Formulae (4.5) and (4.6) were first obtained in Refr. [3].

Remark 1: In the case of inverse scattering in the n -dimensional space one obtains the analogous result as soon as the scattering amplitude is measured.

where

$$\Psi(p, k, \nu) = \int U(x, k) e^{-ip \cdot x} dx,$$

$\delta(p - k\nu)$ represents the incident plane wave, p is a vector in \mathbb{R}^2 , ν is a unit vector in \mathbb{R}^2 and

$$\hat{O}(p) = \int O(x) e^{-ip \cdot x} dx.$$

We introduce a system of coordinates which is related to the direction of propagation of the

initial plane wave. We set

$$p = \eta\nu + \xi\nu^\perp, \quad (5.3)$$

where ν^\perp is the unit vector orthogonal to the vector ν : $\nu = (v_x, v_y)$ and $\nu^\perp = (-v_y, v_x)$.

Let us consider the scattered field $\Psi_{sc}(p, k, \nu) = \Psi(p, k, \nu) - \delta(p - k\nu)$. We find

$$\Psi_{sc}(p, k, \nu) = \frac{k^2}{k^2 - |p|^2 - i0} \int \hat{O}(p - p') \Psi(p', k, \nu) dp', \quad (5.4)$$

In the system of coordinates (5.2) we have

$$O_{LP}(x) = \frac{ik}{4\pi^3} \int_{|\nu|=1} \int_{|\mu|=1} (1-(\mu \cdot \nu)^2)^{1/2} |\mu \cdot \nu| e^{-ik|\mu \cdot \nu|y} \Psi_{sc}^b(y, \mu, k, \nu) e^{i(k\nu - k\mu) \cdot x} d\omega_\nu d\omega_\mu. \quad (5.9)$$

The formula (5.9) is a backpropagation inversion formula which was first obtained by A. J. Devaney [1] and is presented here in a slightly different form.

The case of Rytov approximation is analogous to Born approximation and can be found in Ref. [1]. We note that in the case of a plane incident wave there is a simple relation